*Mycena* sp 21108



Mycena sp 21108 © Pat Leonard

**Cap:** campanulate; up to 3 mms diameter; white with pinkish tone caused by droplets at centre, margin sulcate.

**Stipe:** cylindrical;  $15 - 20 \times 0.25 - 0.5$  mms; hyaline; bulbous hispid ochraceous base.

**Gills:** adnexed, white; 10 reaching stipe. **Flesh:** very thin, white, unchanging.

**Spores:** white; subcylindric;  $3.5 - 4.5 \times 7 - 9 \mu m$ ; Q = 1.9 - 2.1.

Cheilocystidia:

Caulocystidia: narrowly langeniform, some branching in two, smooth.

**Pileipellis:** a cutis of elongated acanthocysts 75 -  $150 \times 12$  -  $20 \mu m$  and disarticulating spaeropedunculate acanthocyst 25 -  $35 \times 20$  -  $30 \mu m$ .

**Habitat:** on leaf in wet sclerophyl forest with *Eucalyptus* 

**Notes:** this minute pink *Mycena* does not key readily in Grgurinovic. It appears to fit most readily in the *Adonideae*, but is not *M. wubabulna* which has a different cap cuticle and caulocystidia.

**Collections examined:** PL 21108, Maroochy Bushland Botanic Garden, Pat Leonard, 9 Nov 08.