Gymnopilus parrumbalus



Gymnopilus parrumbalus © Pat Leonard

Cap: planoconvex with a shallow umbo; 65 - 90 mm diameter; fibrillose at least in outer half; mustard yellow with a darker ochraceous centre (4A7 5B7).

Stipe: cylindrical to slightly tapering downwards, hollow; $30 - 50 \times 6 - 8$ mm; striate fibrillose; yellow ochraceous with a pale zone at the apex.

Gills: adnate, moderately spaced; quite thick; ochraceous orange.

Flesh: ochraceous.

Smell: none.

Spore print: orange-brown.

Spores: ellipsoid; $8 - 11 \times 7 - 7$ µm; finely warty, roughened.

Basidia: clavate; $25 - 35 \times 8 - 10 \mu m$; four-spored.

Cheilocystidia: numerous; fusoid ventricose and many capitate; $25 - 30 \times 5 - 9 \mu m$.

Pleurocystidia: absent.

Pileipellis: a cutis of repent hyphae $5 - 8 \mu m$ wide.

Habitat: in sandy soil with buried wood in Wallum heathland. High woody content resulting from the construction of a firebreak.

Notes: the bright mustard colour with a darker umbo suggests this may be a *Gymnopilus* and this is confirmed by the warty spores. Absence of a veil and spore size confirm the species.

Collections examined: PL300318, Chermside Hills Reserve, Pat Leonard, 18 Mar 2018.