## Gymnopus sp. "Maroochy brown"



Gymnopus sp. "Maroochy brown" © Patrick Leonard

Cap: convex, sometimes shallowly umbonate in the centre; 25 - 80 mm diameter; glabrous, slightly greasy; sulcate striate; light brown, brownish orange (7C4, 7D4), darker brown in the centre on some fruiting bodies; margin irregular.

**Stipe:** cylindrical;  $60 - 80 \times 4 - 5$  mm; glabrous; pale buff with prominent striations that are concolorous with the cap; growing in clusters but not caespitose.

**Gills:** adnexed; moderately crowded; pale buff and a very small paler ring where they join the stipe; with lamellulae in two series.

Flesh: thin; very tough, buff.

Spore print: cream.

**Spores:** ellipsoid;  $10 - 12 \times 4 - 5.5 \mu m$ ; with a prominent oil drop.

**Basidia:** clavate;  $15 - 22 \times 6 - 8 \mu m$ ; 4-spored.

Cheilocystidia: broadly clavate, capitate, some with two lateral or terminal

projections;  $50 - 85 \times 10 - 18 \mu m$ 

**Pleurocystidia:** clavate to fusoid;  $28 - 35 \times 7 - 9 \mu m$ ; with dark granular elements on

the surface.

**Pileipellis:** a cutis of repent hyphae, clamp connections present.

**Substrate:** leaf litter with a high woody content.

**Habitat:** wet sclerophyll forest dominated by *Eucalyptus* species.

**Notes:** a tall and tough collybioid *Gymnopus* with a mid-brown cap and a noticeably

striate stipe. It does not accord with any of the descriptions in Grgurinovic. **Collections examined:** PL80124, Maroochy Bushland Botanic Garden, Patrick

Leonard, 10 Jan 2024.