

Subgenus	Section	Subsection	Series	Type	Species in FoQ
<i>Marasmius</i>	<i>Crinis-eques</i>			<i>M. crinis-equi</i>	<i>crinis-equi</i> , <i>tropicus</i> , <i>kabakada</i> , <i>perumbilicatus</i>
	<i>Marasmius</i>	<i>Marasmius</i>		<i>M. rotula</i>	<i>aff. leucorotalis</i>
		<i>Guyanenses</i>		<i>M. guyanensis</i>	
		<i>Horriduli</i>		<i>M. horridulus</i>	
	<i>Sicciformes</i>	<i>Sicciformes</i>		<i>M. curreyi</i>	<i>ruforotula</i> , <i>nigrobrunneus</i> ,
		<i>Neosessiles</i>		<i>M. neosessilis</i>	<i>tenuissimus</i>
		<i>Leveilleani</i>		<i>M. leveilleanus</i>	<i>leveilleanus</i>
	<i>Variabilispori</i>			<i>M. variabilisporus</i>	<i>tangerinus?</i> (probably not) 'dimpled buff orange', 'orange cushion'
	<i>Sanguirotales</i>			<i>M. sanguirotalis</i>	<i>argillaceus?</i> (probably not)
<i>Globulares</i>	<i>Globulares</i>	<i>Leonini</i>	<i>Corrugati</i>	<i>M. corrugatus</i>	<i>pellucidus</i>
			<i>Graminicolae</i>	<i>M. graminicola</i>	<i>suthepensis</i> , <i>koae</i> , <i>Clav. extendens</i> (maybe?)
			<i>Bambusiniiformis</i>	<i>M. bambusiniiformis</i>	<i>clocca</i>
			<i>Leonini</i>	<i>M. leoninus</i>	
			<i>Multicystidiati</i>	<i>M. multicystidiatus</i>	<i>multicystidiatus</i>
			<i>Luteoli</i>	<i>M. luteolus</i>	<i>elegans</i> , <i>pseudoelegans</i> , <i>dilkusha</i> , <i>leppii</i>
		<i>Fusicystides</i>		<i>M. isabellinus</i>	<i>brunneolorobustus?</i>
		<i>Atrorubentes</i>	<i>Atrorubentes</i>	<i>M. atrorubens</i>	<i>paratrichotus</i> , <i>croceus</i> , <i>australotrichotus</i>
			<i>Conferti</i>	<i>M. confertus</i>	
			<i>Crinipes</i>	<i>M. crinipes</i>	<i>lebeliae</i>
			<i>Hinnulei</i>	<i>M. hinnuleus</i>	
			<i>Imitarii</i>	<i>M. imitarius</i>	
			<i>Pulcherripes</i>	<i>M. pulcherripes</i>	
			<i>Purpureostriati</i>	<i>M. purpureostriatus</i>	<i>musisporus</i> , <i>carbinensis</i> , <i>ballator</i> , 'pink stripe <i>Toowoomba</i> '
			<i>Rhabarbarini</i>	<i>M. rhabarbarinus</i>	
		<i>Globulares</i>	<i>Wynnearum</i>	<i>M. wynneae</i>	<i>oreades</i>
			<i>Brunneospermi</i>	<i>M. brunneospermus</i>	
			<i>Cohaerentes</i>	<i>M. cohaerens</i>	
			<i>Silvicolae</i>	<i>M. silvicola</i>	
			<i>Hypochroides</i>	<i>M. hypochroides</i>	<i>vagus</i>
	<i>Sicci</i>		<i>Ferruginei</i>	<i>M. ferrugineus</i>	
			<i>Sicci</i>	<i>M. siccus</i>	
			<i>Haematocephali</i>	<i>M. haematocephali</i>	<i>wunga</i> , <i>wianwian</i> , 'clay pink' <i>campaniformis</i> , <i>vinaceus</i> , <i>purpureospadiceus</i> , <i>porphyrocephalus</i> , <i>pervagatus</i>

This Table follows the classification of *Marasmius* proposed by Oliveira et al (2024) in **Taxonomic revision of *Marasmius* (Fr.) and *Marasmiaceae* (Roze ex Kühner) based on multigene phylogenetics and morphological evidence** in *Fungal Diversity*, **127**, 1–54. Sections, subsections or series that have not been found to date in Australia are in brown. All type species are in purple. Species with “probably not” noted against them await further clarification.